



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 911 399 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
28.04.1999 Bulletin 1999/17

(21) Application number: 98203240.1

(22) Date of filing: 25.09.1998

(51) Int. Cl.⁶: **C12N 15/12**, **C07K 14/47**,
C07K 16/18, **A61K 48/00**,
A61K 31/70, **C12Q 1/68**,
G01N 33/68, **C12N 1/21**,
C12N 5/10

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **01.10.1997 US 60623 P**
16.03.1998 US 39609

(71) Applicant:
SMITHKLINE BEECHAM CORPORATION
Philadelphia Pennsylvania 19103 (US)

(72) Inventors:
• **Albone, Earl Francis**
King of Prussia, PA 19406 (US)
• **Kikly, Kristine Kay**
King of Prussia, PA 19406 (US)

(74) Representative:
Crump, Julian Richard John et al
FJ Cleveland,
40-43 Chancery Lane
London WC2A 1JQ (GB)

(54) **A kringle-related clone, HTHBZ47**

(57) HTHBZ47 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing HTHBZ47 polypeptides and polynucleotides in therapy, and diagnostic assays for such.

EP 0 911 399 A2

Description

[0001] This patent claims the benefit of U.S. Provisional Application No. 60/060,623, filed October 1, 1997, whose contents are incorporated herein by reference.

Field of the Invention

[0002] This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides, to their use in therapy and in identifying compounds which may be agonists, antagonists and/or inhibitors which are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

Background of the Invention

[0003] The drug discovery process is currently undergoing a fundamental revolution as it embraces 'functional genomics', that is, high throughput genome- or gene-based biology. This approach is rapidly superceding earlier approaches based on 'positional cloning'. A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

[0004] Functional genomics relies heavily on the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

Summary of the Invention

[0005] The present invention relates to HTHBZ47, in particular HTHBZ47 polypeptides and HTHBZ47 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including the treatment of cancer, inflammation, autoimmunity, allergy, asthma, rheumatoid arthritis, CNS inflammation, cerebellar degeneration, Alzheimer's disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, head injury damage, and other neurological abnormalities, septic shock, sepsis, stroke, osteoporosis, osteoarthritis, ischemia reperfusion injury, cardiovascular disease, kidney disease, liver disease, ischemic injury, myocardial infarction, hypotension, hypertension, AIDS, myelodysplastic syndromes and other hematologic abnormalities, aplastic anemia, male pattern baldness, and bacterial, fungal, protozoan and viral infections, hereinafter referred to as "the Diseases", amongst others. In a further aspect, the invention relates to methods for identifying agonists and antagonists/inhibitors using the materials provided by the invention, and treating conditions associated with HTHBZ47 imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate HTHBZ47 activity or levels.

Description of the Invention

[0006] In a first aspect, the present invention relates to HTHBZ47 polypeptides. Such peptides include isolated polypeptides comprising an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2 over the entire length of SEQ ID NO:2. Such polypeptides include those comprising the amino acid of SEQ ID NO:2.

[0007] Further peptides of the present invention include isolated polypeptides in which the amino acid sequence has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2. Such polypeptides include the polypeptide of SEQ ID NO:2.

[0008] Further peptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:1.

[0009] Polypeptides of the present invention are members of the kringle family of polypeptides. They are therefore of interest because kringle domains are found in proteins involved in blood coagulation (thrombin, factor XII), fibrinolysis (tissue plasminogen activator, urokinase), liver regeneration (hepatocyte growth factor), and lipid metabolism (apolipoprotein A). These proteins may have a single kringle domain, as in urokinase, or multiple copies, as in apolipoprotein A, which has 38 repeated kringle domains. Kringle motif-containing proteins often have other separate motifs which perform different functions, such as growth factor and serine protease domains (Campbell, I. D. et al, Philos. Trans. R. Soc. Lon. B. Biol. Sci., 332(1263), p. 165-170, 1991). Kringle domains are thought to aid in binding to fibrin clots or membrane components, as well as to modulate protease activity in proteins having this activity associated with them (Lijnen, H. R. et al, Ballin's Clin. Hematol., 8(2), p. 277-290, 1995).

[0010] The proteins, tissue plasminogen activator (t-PA) and urokinase, have both been used clinically in the treatment of coronary heart disease in both its acute and chronic phases (Gulba, D. C. et al, Ann. Hematol., 73 Suppl 1, p. S9-27, 1996), and t-PA has been used clinically in the management of ischemic stroke (Albers, G. W., West J Med., 166(4), p. 253-62, 1994). Both of these proteins aid in the course of the disease by binding to fibrin clots and activating plasminogen, which degrades the clot by proteolysis. Other kringle-containing proteins, such as hepatocyte growth factor, have been implicated in metastatic cancers, presumably via growth factor domains associated with these proteins (Weidner, K. M. et al, EXS, 65, p. 311-28, 1993). These properties are hereinafter referred to as "HTHBZ47 activity" or "HTHBZ47 polypeptide activity" or "biological activity of HTHBZ47". Also included amongst these activities are antigenic and immunogenic activities of said HTHBZ47 polypeptides, in particular the antigenic and immunogenic activities of the polypeptide of SEQ ID NO:2. Preferably, a polypeptide of the present invention exhibits at least one biological activity of HTHBZ47.

[0011] The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

[0012] The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

[0013] Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

[0014] In a further aspect, the present invention relates to HTHBZ47 polynucleotides. Such polynucleotides include isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ ID NO:2. In this regard, polypeptides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:1 encoding the polypeptide of SEQ ID NO:2.

[0015] Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence that has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2, over the entire coding region. In this regard, polynucleotides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred.

[0016] Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to SEQ ID NO:1 over the entire length of SEQ ID NO:1. In this regard, polynucleotides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the polynucleotide of SEQ ID NO:1 as well as the polynucleotide of SEQ ID NO:1.

[0017] The invention also provides polynucleotides which are complementary to all the above described polynucleotides.

[0018] The nucleotide sequence of SEQ ID NO:1 shows homology with human tissue-type plasminogen activator (Harris, T. J. et al., Mol. Biol. Med. 3(3) : 279-292, 1986), Accession # M15518. The nucleotide sequence of SEQ ID NO:1 is a cDNA sequence and comprises a polypeptide encoding sequence (nucleotide 87 to 1463) encoding a polypeptide of 458 amino acids, the polypeptide of SEQ ID NO:2. The nucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in SEQ ID NO:1 or it may be a sequence other than the one contained in SEQ ID NO:1, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2. The polypeptide of SEQ ID NO:2 is structurally related to other proteins of the kringle family, having homology and/or structural similarity with tissue plasminogen activator precursor (tPA), (Harris et al., Mol. Biol. Med. 3(3) : 279-292, 1986, GenBank accession # 137119).

[0019] Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one HTHBZ47 activity.

[0020] The present invention also relates to partial or other polynucleotide and polypeptide sequences which were

first identified prior to the determination of the corresponding full length sequences of SEQ ID NO:1 and SEQ ID NO:2.

[0021] Accordingly, in a further aspect, the present invention provides for an isolated polynucleotide comprising:

- (a) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;
- (b) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity, to SEQ ID NO:1 over the entire length of SEQ ID NO:3;
- (c) the polynucleotide of SEQ ID NO:3; or
- (d) a nucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:4, over the entire length of SEQ ID NO:4, as well as the polynucleotide of SEQ ID NO:3.

[0022] The present invention further provides for a polypeptide which:

- (a) comprises an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2 over the entire length of SEQ ID NO:4;
 - (b) has an amino acid sequence which is at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:4;
 - (c) comprises the amino acid of SEQ ID NO:4; and
 - (d) is the polypeptide of SEQ ID NO:4;
- as well as polypeptides encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:3.

[0023] The nucleotide sequence of SEQ ID NO:3 and the peptide sequence encoded thereby are derived from EST (Expressed Sequence Tag) sequences. It is recognised by those skilled in the art that there will inevitably be some nucleotide sequence reading errors in EST sequences (see Adams, M.D. *et al.*, Nature 377 (supp) 3, 1995). Accordingly, the nucleotide sequence of SEQ ID NO:3 and the peptide sequence encoded therefrom are therefore subject to the same inherent limitations in sequence accuracy. Furthermore, the peptide sequence encoded by SEQ ID NO:3 comprises a region of identity or close homology and/or close structural similarity (for example a conservative amino acid difference) with the closest homologous or structurally similar protein.

[0024] Polynucleotides of the present invention may be obtained, using standard cloning and screening techniques, from a cDNA library derived from mRNA in cells of human thymus and fetal liver, using the expressed sequence tag (EST) analysis (Adams, M.D., *et al.* Science (1991) 252:1651-1656; Adams, M.D. *et al.*, Nature, (1992) 355:632-634; Adams, M.D., *et al.*, Nature (1995) 377 Supp.3:174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

[0025] When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself; or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

[0026] Further embodiments of the present invention include polynucleotides encoding polypeptide variants which comprise the amino acid sequence of SEQ ID NO:2 and in which several, for instance from 5 to 10, 1 to 5, 1 to 3, 1 to 2 or 1, amino acid residues are substituted, deleted or added, in any combination.

[0027] Polynucleotides which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:1, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification (PCR) reaction, to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs from species other than human) that have a high sequence similarity to SEQ ID NO:1. Typically these nucleotide sequences are 70% identical, preferably 80% identical, more preferably 90% identical, most preferably 95% identical to that of the referent. The probes or primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have

at least 50 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides.

[0028] A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than human, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof, and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes polynucleotides obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof.

[0029] The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide is cut short at the 5' end of the cDNA. This is a consequence of reverse transcriptase, an enzyme with inherently low 'processivity' (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during 1st strand cDNA synthesis.

[0030] There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., PNAS USA 85: 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the 'missing' 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analysed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

[0031] Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems which comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

[0032] For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., Basic Methods in Molecular Biology (1986) and Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). Preferred such methods include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

[0033] Representative examples of appropriate hosts include bacterial cells, such as *streptococci*, *staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

[0034] A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector which is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (*supra*). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

[0035] If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

5 [0036] Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

10 [0037] This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO:1 which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

15 [0038] Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled HTHBZ47 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, e.g., Myers *et al.*, *Science* (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, *Proc Natl Acad Sci USA* (1985) 85: 4397-4401). In another embodiment, an array of oligonucleotide probes comprising HTHBZ47 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

30 [0039] The diagnostic assays offer a process for diagnosing or determining a susceptibility to the Diseases through detection of mutation in the HTHBZ47 gene by the methods described. In addition, such diseases may be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of polypeptide or mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

40 [0040] Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment thereof;

(b) a nucleotide sequence complementary to that of (a);

45 (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

[0041] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly cancer, inflammation, autoimmunity, allergy, asthma, rheumatoid arthritis, CNS inflammation, cerebellar degeneration, Alzheimer's disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, head injury damage, and other neurological abnormalities, septic shock, sepsis, stroke, osteoporosis, osteoarthritis, ischemia reperfusion injury, cardiovascular disease, kidney disease, liver disease, ischemic injury, myocardial infarction, hypotension, hypertension, AIDS, myelodysplastic syndromes and other hematologic abnormalities, aplastic anemia, male pattern baldness, and bacterial, fungal, protozoan and viral infections, amongst others.

55 [0042] The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in cor-

relating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

[0043] The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

[0044] The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them, can also be used as immunogens to produce antibodies immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

[0045] Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

[0046] Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

[0047] The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

[0048] Antibodies against polypeptides of the present invention may also be employed to treat the Diseases, amongst others.

[0049] In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

[0050] Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response to protect said animal from the Diseases hereinbefore mentioned, amongst others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering a polypeptide of the present invention via a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

[0051] A further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a polypeptide of the present invention wherein the composition comprises a polypeptide or polynucleotide of the present invention. The vaccine formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

[0052] Polypeptides of the present invention are responsible for many biological functions, including many disease states, in particular the Diseases hereinbefore mentioned. It is therefore desirable to devise screening methods to identify compounds which stimulate or which inhibit the function of the polypeptide. Accordingly, in a further aspect, the

present invention provides for a method of screening compounds to identify those which stimulate or which inhibit the function of the polypeptide. In general, agonists or antagonists may be employed for therapeutic and prophylactic purposes for such Diseases as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. Such agonists, antagonists or inhibitors so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; or may be structural or functional mimetics thereof (see Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991)).

[0053] The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring HTHBZ47 activity in the mixture, and comparing the HTHBZ47 activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and HTHBZ47 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, *J Mol Recognition*, 8:52-58 (1995); and K. Johanson *et al.*, *J Biol Chem*, 270(16):9459-9471 (1995)).

[0054] The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

[0055] The polypeptide may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance, ^{125}I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide which compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

[0056] Examples of potential polypeptide antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or small molecules which bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

[0057] Thus, in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for polypeptides of the present invention; or compounds which decrease or enhance the production of such polypeptides, which comprises:

- (a) a polypeptide of the present invention;
 - (b) a recombinant cell expressing a polypeptide of the present invention;
 - (c) a cell membrane expressing a polypeptide of the present invention; or
 - (d) antibody to a polypeptide of the present invention;
- which polypeptide is preferably that of SEQ ID NO:2.

[0058] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

[0059] It will be readily appreciated by the skilled artisan that a polypeptide of the present invention may also be used in a method for the structure-based design of an agonist, antagonist or inhibitor of the polypeptide, by:

- (a) determining in the first instance the three-dimensional structure of the polypeptide;
- (b) deducing the three-dimensional structure for the likely reactive or binding site(s) of an agonist, antagonist or inhibitor;
- (c) synthesizing candidate compounds that are predicted to bind to or react with the deduced binding or reactive site; and

(d) testing whether the candidate compounds are indeed agonists, antagonists or inhibitors.

It will be further appreciated that this will normally be an interactive process.

[0060] In a further aspect, the present invention provides methods of treating abnormal conditions such as, for instance, cancer, inflammation, autoimmunity, allergy, asthma, rheumatoid arthritis, CNS inflammation, cerebellar degeneration, Alzheimer's disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, head injury damage, and other neurological abnormalities, septic shock, sepsis, stroke, osteoporosis, osteoarthritis, ischemia reperfusion injury, cardiovascular disease, kidney disease, liver disease, ischemic injury, myocardial infarction, hypotension, hypertension, AIDS, myelodysplastic syndromes and other hematologic abnormalities, aplastic anemia, male pattern baldness, and bacterial, fungal, protozoan and viral infections, related to either an excess of, or an under-expression of, HTHBZ47 polypeptide activity.

[0061] If the activity of the polypeptide is in excess, several approaches are available. One approach comprises administering to a subject in need thereof an inhibitor compound (antagonist) as hereinabove described, optionally in combination with a pharmaceutically acceptable carrier, in an amount effective to inhibit the function of the polypeptide, such as, for example, by blocking the binding of ligands, substrates, receptors, enzymes, etc., or by inhibiting a second signal, and thereby alleviating the abnormal condition. In another approach, soluble forms of the polypeptides still capable of binding the ligand, substrate, enzymes, receptors, etc. in competition with endogenous polypeptide may be administered. Typical examples of such competitors include fragments of the HTHBZ47 polypeptide.

[0062] In still another approach, expression of the gene encoding endogenous HTHBZ47 polypeptide can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered (see, for example, O'Connor, *J Neurochem* (1991) 56:560 in *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices with the gene can be supplied (see, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360). These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*.

[0063] For treating abnormal conditions related to an under-expression of HTHBZ47 and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates a polypeptide of the present invention, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of HTHBZ47 by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For an overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in *Human Molecular Genetics*, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of a polypeptide of the present invention in combination with a suitable pharmaceutical carrier.

[0064] In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide, such as the soluble form of a polypeptide of the present invention, agonist/antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0065] The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, and the like.

[0066] The dosage range required depends on the choice of peptide or other compounds of the present invention, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies

of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

[0067] Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

[0068] Polynucleotide and polypeptide sequences form a valuable information resource with which to identify further sequences of similar homology. This is most easily facilitated by storing the sequence in a computer readable medium and then using the stored data to search a sequence database using well known searching tools, such as GCC. Accordingly, in a further aspect, the present invention provides for a computer readable medium having stored thereon a polynucleotide comprising the sequence of SEQ ID NO:1 and/or a polypeptide sequence encoded thereby.

[0069] The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

[0070] "Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

[0071] "Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated," as the term is employed herein.

[0072] "Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide; which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

[0073] "Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., Post-translational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62).

[0074] "Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, refer-

enc polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

[0075] "Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" *per se* has an art-recognized meaning and can be calculated using published techniques (see, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., *Nucleic Acids Research* (1984) 12(1):387), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J Molec Biol* (1990) 215:403).

[0076] By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is be 100% identical, or it may include up to a certain integer number of nucleotide alterations as compared to the reference sequence. Such alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the numerical percent of the respective percent identity (divided by 100) and subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y),$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, and y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

[0077] Similarly, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the % identity is less than 100%. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the numerical percent of the respective percent identity (divided by 100) and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, and y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

[0078] "Fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0 464 discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, employing an immunoglobulin-Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties [see, e.g., EP-A 0232 262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified.

[0079] All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

15

20

25

30

35

40

45

50

55

SEQUENCE INFORMATION

SEQ ID NO:1

5 CTGCGCCTGCTCCCATGGCCGCCCGGCTCCCGCGCTGCCCCCTTTACCCCGGGCCGCGCCCGGGGCCCGGC
 ACTGACGGCCCATGGCGCCGCCAGCCGCCCGCTCGCCCTGCTCTCCGCGCGGGCGCTCAGCTGGCGGCCCGGC
 CCGCGCCTAGCCCGGCTCGGCCCGGACCCGAGTGTTCACAGCCAATGGTGGGATTATAGGGGAACACAGA
 10 ACTGGACAGCACTACAAGGCGGGAAGCCATGTCTGTTTTGGAACGAGACTTTCAGCATCCATACAACTCTGA
 AATACCCCAACGGTAGTGGGGGCTGGGTGAGCACAATATTGCAGAAATCCAGATGGAG
 ACGTGAGCCCTGGTGCTATGTGGCAGAGCAGAGGATGGTGTCTACTGGAAGTACTGTGAGATACCTGCTTGCC
 AGATGCTGGAAACCTTGGCTGCTACAAGGATCATGGAACCCACCTCCTCTAACTGGCACCAGTAAACGCTCA
 ACAAACTCACCATACAAATTCATCAGTTTTTGTGGAGTCAGAGGTTCAAGTTTGTGGGATGGAGTCAGGCT
 15 ATGCTTGCTTCTGTGGAACAATCTGATTACTGGAAGTACGGGGAGGCAGCCAGTACCGAATGCAACAGCGTCT
 GCTTCGGGGATCACACCAACCTGTGGTGGCGATGGCAGGATCATCCTCTTTGATACTC
 TCGTGGGCGCTGCGGTGGGAATCTCAGCCATGTCTTCTGTGGTCTATTCCCTGACTTCCCCGACACCTATG
 CCACGGGGAGGGTCTGCTACTGGACCATCCGGGTTCCGGGGGCTCCCATCCACTTCAGCTTCCCCCTATTTG
 20 ACATCAGGGACTCGGCGGACATGGTGGAGCTTCTGGATGGCTACACCCACCGTGTCTAGCCCGCTTCCACGGGA
 GGAGCCGCCACCTCTGTCTTCAACGTCTCTGTGACTTCTGTCATCTTGTATTCTTCTCTGATCGCATCAATC
 AGGCCAGGGATTGTCTGTTTTATACCAAGCCGTCAAGGAAGAACTGCCACAGGAGAGGC
 CCGCTGTCAACCAGACGGTGGCCGAGGTGATCAGGAGCAGGCCAACCTCAGTGTGAGCGCTGCCCGTCTCCCA
 AAGTCTCTATGTATCACCACAGCCCCAGCCACCCACCTCAAAGTGTCCAGAATGGACAGTCTATGGTCTGG
 25 CAACTCTCTCATCTCACAGTCACAGCCATTGTAGCAAGATACTTCTGCACGTACATTCAAATCCCATCGTG
 TTCTGCTTCAGGGACCTTAGGGATTGTATCAACCAGGACTTCGGGGGAAATCTGGAGCATTTTTTTACAAGC
 CTTCCACTTCAATTTCCATCTTTAAGAAGAACTCAAGGTCAGAGTCAACAAGATGACC
 GCAATCCCCTTGTGAGTGAATAAAACCCACTGTGCTAGGACTTGAGGTCCCTCTTTGAGCTCAAGGCTGCCG
 30 TGGTCAACCTCTCTGTGGTCTTCTCTGACAGACTTCCCCCTCTCTCCCTCTGCTCGGCTCTTCCGGGAA
 CCTCTCTACAGACTAGGAAGAGGCACCTGTGCCAGGGCAGGCAGAGCTGGATTCTCTCTGCTTATCGATT
 GCACCTTAGGAGAGAGACTCAAAGCCCTGGGCCCGGCTCTCTGATCTCTCTGATCTAGCTAGCAGTGGGGG
 TGTCAGGACAGTGAGGCTGAGATGACAGAGGTGGTATGGCTGGCACAGGGCTCAGGTAC
 35 ATTCTAGATGGCTGTGAGGTGGTGGTAGCTTTAGTTACATTGAATTTTCTTGCTTCTATTTTGTCCACAC

SEQ ID NO:2

40 MAPPAARLALLSAAALTLAARPAAPSPGLGPGPECFTANGADYRGTQNWALQGGKPCLPWNETFOHPYNTLKYPN
 GSGGLGEHNYCRNPDGDVSPWCYVAHEDGVYWKYCEIPACQMPGNLGCYKDHGNPPPLTGTSKTSNKLTIQTCI
 SFCRSQRFKPFAGMESGYACFCGNNPDYWKYGEAASTECSVCFGDHTQPCGGDGRIILFDLTVGACGGNYSAMSS
 VVYSPDFPDYATGRVCYWTIRVPGASHIHFSFPLEDIRDSADMVELLDGYTHRVLARFHGRSRPPLSPNVSLDF
 45 VILYFFSDRINQAQGFVLYQAVKEELPQERPAVNQTVAEVITEQANLSVSAARSSKVLV
 VITTSFSPHPQTVPETVYGLATLLILTVTAIVAKILLHVTFKSHRVPASGDLRDCHQPGTSGEINSIFYKPSTS
 ISIFKQKLKGSQDDRNPLVSD

SEQ ID NO:3

50 GGCACGAGCTGGACAGCACTACAAGCGGGGAAGCCATGTCTGTTTTGGAACGAGACTTTCAGCATCCATACAAC
 ACTCTGAAATACCCCAACGGTAGTGGGGGCTGGGTGAGCACAATATTGCAGAAATCCAGATGGAGACGTGAGC
 CCCTGGTGCTATGTGGCAGAGCAGGAGGATGGTGTCTACTGGAAGTACTGTGAGATACCTGCTTGCCAGATGCCT
 GGAAACCTTGGCTGCTACAAGGATCATGGAACCCACCTCCTCTAACTGGCACCATAAAACGTCACAACTC

5 ACCATACAAACTTGCATCAGTTTTTGTGCGAGTCAGAGGTTCAAGTTTGCTGGGATGGAGTCAGGCTATGCTTGC
TTCTGTGGAAACAATCCTGATTACTGGAAGTACGGGGAGGCAGCCAGTACCGAATGCAACAGCGTCTGCTTCGGG
GATCACACCCAACCTGTGGTGGCGATGGCAGGATCATCTCTTTGATACTCTCGTGGGCGCCTGCGGTGGGAAC
TACTCAGCCATGTCTTCTGTGGTCTATTCCCCTGACTTCCCCGACACCTATGCCACGGGGAGGGTCTGCTACTGG
ACCATCCGGGTTCGGGGGGCCTCCACATCCACTTCAGCTTCCCCCTATTTGACATCAGGGACTCGGCGGACATG
GTGGAGCTTCTGGATGGCTACACCCACCGTGTCTTAGCCCGCTTCACGGGAGGAGCCGCCACCTCTGTCTTC
10 AACGTCTCTCTGGACTTCGTTCATCTTGTATTTCTTCTCTGATCGCATCAATCAGGCCCAGGGATTGCTGTTTTA
TACCAAGCGGTCAAGGAAGAACTGCCACAGGAGAGGCCCGCTGTCAACCAGACGGTGGCCGAGGTGGATCACGGA
GCAGGCCAACCTTCAGTGTTCAGCGCTTCCCGGTCTCCTCAAAGTCTATTATGTGCATGCACCA

SEQ ID NO:4

15 GTSWTALQGG KPCLFWNETF QHPYNTLKYP NGSGGLGEHN YCRNPDGDVS PWCYVAEHED
GVYWKYCEIP ACOMPONLGC YKDHGNPPPL TGTIKTSNKL TIQTCISFCR SQRKFAGME
SGYACFCGNN PDYWKYGEAA STECNSVCFG DHTQPCGGDG RIILFDTLVG ACGGNYSAMS
SVVYSPDFPD TYATGRVCYW TIRVPGASHI HFSFPLFDIR DSADMVELLD GYTHRVLARF
HGRSRPPLSF NVSLDFVILY FFSDRINQAO GFAVLYQAVK BELPQERPAV NQTVAEVDHG
AGQPSVFSAS RSPQSPIMCM H

20

25

30

35

40

45

50

55

SEQUENCE LISTING

5 (1) GENERAL INFORMATION

(i) APPLICANT

(A) NAME: SMITHKLINE BEECHAM CORPORATION
10 (B) STREET: ONE FRANKLIN PLAZA
(C) CITY: PHILADELPHIA
(D) STATE OR PROVINCE: PENNSYLVANIA
(E) COUNTRY: USA
15 (F) POSTAL CODE: 19103

(ii) TITLE OF THE INVENTION: A KRINGLE-RELATED CLONE,
HTHB247

20 (iii) NUMBER OF SEQUENCES: 4

(iv) COMPUTER-READABLE FORM:

(A) MEDIUM TYPE: Diskette
25 (B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: DOS
(D) SOFTWARE: FastSEQ for Windows Version 2.0

30 (v) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: TO BE ASSIGNED

35 (2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1875 base pairs
(B) TYPE: nucleic acid
40 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCGCCCTGC TCCCATGGCC GCCCCCGGCT CCCC GCGCTG CCCCCTTTAC CCCGGGCCGC 60
50 GCCCCGGGGC CCCGCACTGA CGGCCCATGG CGCCGCCAGC CGCCCGCCTC GCCCTGCTCT 120
CCGCCCGGGC GCTCACGCTG GCGGCCCGGC CCGCGCCTAG CCCC GGCCCTC GGCCCCGGAC 180
CCGAGTGTTT CACAGCCAAT GGTGCGGATT ATAGGGGAAC ACAGAACTGG ACAGCACTAC 240

55

EP 0 911 399 A2

5
10
15
20
25
30
35

AAGGCGGGAA GCCATGTCTG TTTTGGAAAG AGACTTTCCA GCATCCATAC AACACTCTGA 300
AATACCCCAA CGGTAGTGGG GGCTGGGTG AGCACACTA TTGCAGAAAT CCAGATGGAG 360
ACGTGAGCCC CTGGTGCTAT GTGGCAGAGC ACGAGGATGG TGTCTACTGG AAGTACTGTG 420
AGATACCTGC TTGCCAGATG CCTGGAAACC TTGGCTGCTA CAAGGATCAT GGAAACCCAC 480
CTCCTCTAAC TGGCACCAGT AAAACGTCCA ACAAACCTCAC CATACAAAC TGCATCAGTT 540
TTTGTCGGAG TCAGAGGTTC AAGTTTGTG GGATGGAGTC AGGCTATGCT TGCTTCTGTG 600
GAAACAATCC TGATTACTGG AAGTACGGGG AGGCAGCCAG TACCGAATGC AACAGCGTCT 660
GCTTCGGGGA TCACACCCAA CCCTGTGGTG GCGATGGCAG GATCATCTC TTTGATACTC 720
TCGTGGGCGC CTGCGGTGGG AACTACTCAG CCATGTCTTC TGTGGTCTAT TCCCCTGACT 780
TCCCCGACAC CTATGCCACG GGGAGGTCT GCTACTGGAC CATCCGGGT CCGGGGGCCT 840
CCCACATCCA CTTAGCTTC CCCCTATTG ACATCAGGA CTCGGCGGAC ATGGTGGAGC 900
TTCTGGATGG CTACACCCAC CGTGTCTAG CCCGCTTCCA CGGGAGGAGC CGCCACCTC 960
TGTCCTTCAA CGTCTCTCTG GACTTCGTCA TCTTGTATT CTTCTCTGAT CGCATCAATC 1020
AGGCCAGGG ATTGCTGTT TTATACCAAG CCGTCAAGGA AGAACTGCCA CAGGAGAGGC 1080
CCGCTGTCAA CCAGACGGTG GCCGAGGTGA TCACGGAGCA GGCCAACCTC AGTGTACAGC 1140
CTGCCCGTCT CTTCAAAGTC CTCTATGTCA TCACCACCG CCCCAGCCAC CCACCTCAAA 1200
CTGTCCCAGA ATGSACAGTC TATGGTCTGG CAACTCTCT CATCTCACA GTCACAGCCA 1260
TTGTAGCAAA GATACTTCTG CACGTCACAT TCAAATCCCA TCGTGTTCCT GCTTCAGGGG 1320
ACCTTAGGGA TTGTATCAA CCAGGGACTT CGGGGAAAT CTGGAGCATT TTTTACAAGC 1380
CTTCCACTTC AATTTCATC TTTAAGAAGA AACTCAAGGG TCAGAGTCAA CAAGATGACC 1440
GCAATCCCCT TGTGAGTGAC TAAAAACCC ACTGTGCCTA GGAATGAGG TCCCTCTTGT 1500
AGCTCAAGGC TGGCGTGGTC AACCTCTCT GTGGTTCTT TCTGACAGAC TCTTCCCTC 1560
CTCTCCCTCT GCCTCGGCTC TTCGGGAAA CCTCCTCCTA CAGACTAGGA AGAGGCACCC 1620
TGCTGCCAGG GCAGGCAGAG CTGGATTCTT CTGCTTCAT CGATTGCACT TAGGAGAGAG 1680
ACTCAAAGCC CTGGGCCCGG CCTCTCTGC ATCTCTCTT GATCTAGCTA GCAGTGGGGG 1740
TGTCAGGACA GTGAGGCTGA GATGACAGAG GTGGTCATGG CTGGCACAGG GCTCAGGTAC 1800
ATTCTAGATG GCTGTCAGGT GGTGGGTAGC TTTAGTTACA TTGAATTTT CTTGCTTCTC 1860
TATTTTGTG CACAC 1875

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 458 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Pro Pro Ala Ala Arg Leu Ala Leu Leu Ser Ala Ala Ala Leu
1 5 10 15

EP 0 911 399 A2

	Thr	Leu	Ala	Ala	Arg	Pro	Ala	Pro	Ser	Pro	Gly	Leu	Gly	Pro	Gly	Pro
				20					25					30		
5	Glu	Cys	Phe	Thr	Ala	Asn	Gly	Ala	Asp	Tyr	Arg	Gly	Thr	Gln	Asn	Trp
			35					40					45			
	Thr	Ala	Leu	Gln	Gly	Gly	Lys	Pro	Cys	Leu	Phe	Trp	Asn	Glu	Thr	Phe
		50					55					60				
10	Gln	His	Pro	Tyr	Asn	Thr	Leu	Lys	Tyr	Pro	Asn	Gly	Ser	Gly	Gly	Leu
	65					70					75				80	
	Gly	Glu	His	Asn	Tyr	Cys	Arg	Asn	Pro	Asp	Gly	Asp	Val	Ser	Pro	Trp
				85						90					95	
15	Cys	Tyr	Val	Ala	Glu	His	Glu	Asp	Gly	Val	Tyr	Trp	Lys	Tyr	Cys	Glu
				100					105						110	
	Ile	Pro	Ala	Cys	Gln	Met	Pro	Gly	Asn	Leu	Gly	Cys	Tyr	Lys	Asp	His
			115						120					125		
20	Gly	Asn	Pro	Pro	Pro	Leu	Thr	Gly	Thr	Ser	Lys	Thr	Ser	Asn	Lys	Leu
		130					135						140			
	Thr	Ile	Gln	Thr	Cys	Ile	Ser	Phe	Cys	Arg	Ser	Gln	Arg	Phe	Lys	Phe
	145					150					155				160	
25	Ala	Gly	Met	Glu	Ser	Gly	Tyr	Ala	Cys	Phe	Cys	Gly	Asn	Asn	Pro	Asp
				165						170					175	
	Tyr	Trp	Lys	Tyr	Gly	Glu	Ala	Ala	Ser	Thr	Glu	Cys	Asn	Ser	Val	Cys
				180					185					190		
30	Phe	Gly	Asp	His	Thr	Gln	Pro	Cys	Gly	Gly	Asp	Gly	Arg	Ile	Ile	Leu
		195						200					205			
	Phe	Asp	Thr	Leu	Val	Gly	Ala	Cys	Gly	Gly	Asn	Tyr	Ser	Ala	Met	Ser
		210					215					220				
35	Ser	Val	Val	Tyr	Ser	Pro	Asp	Phe	Pro	Asp	Thr	Tyr	Ala	Thr	Gly	Arg
	225					230				235					240	
	Val	Cys	Tyr	Trp	Thr	Ile	Arg	Val	Pro	Gly	Ala	Ser	His	Ile	His	Phe
				245						250					255	
40	Ser	Phe	Pro	Leu	Phe	Asp	Ile	Arg	Asp	Ser	Ala	Asp	Met	Val	Glu	Leu
			260						265					270		
	Leu	Asp	Gly	Tyr	Thr	His	Arg	Val	Leu	Ala	Arg	Phe	His	Gly	Arg	Ser
		275						280					285			
45	Arg	Pro	Pro	Leu	Ser	Phe	Asn	Val	Ser	Leu	Asp	Phe	Val	Ile	Leu	Tyr
		290					295					300				
	Phe	Phe	Ser	Asp	Arg	Ile	Asn	Gln	Ala	Gln	Gly	Phe	Ala	Val	Leu	Tyr
	305					310					315				320	
	Gln	Ala	Val	Lys	Glu	Glu	Leu	Pro	Gln	Glu	Arg	Pro	Ala	Val	Asn	Gln
				325						330					335	
50	Thr	Val	Ala	Glu	Val	Ile	Thr	Glu	Gln	Ala	Asn	Leu	Ser	Val	Ser	Ala
				340					345					350		
	Ala	Arg	Ser	Ser	Lys	Val	Leu	Tyr	Val	Ile	Thr	Thr	Ser	Pro	Ser	His

355 360 365
 Pro Pro Gln Thr Val Pro Glu Trp Thr Val Tyr Gly Leu Ala Thr Leu
 5 370 375 380
 Leu Ile Leu Thr Val Thr Ala Ile Val Ala Lys Ile Leu Leu His Val
 385 390 395 400
 Thr Phe Lys Ser His Arg Val Pro Ala Ser Gly Asp Leu Arg Asp Cys
 405 410 415
 10 His Gln Pro Gly Thr Ser Gly Glu Ile Trp Ser Ile Phe Tyr Lys Pro
 420 425 430
 Ser Thr Ser Ile Ser Ile Phe Lys Lys Lys Leu Lys Gly Gln Ser Gln
 435 440 445
 15 Gln Asp Asp Arg Asn Pro Leu Val Ser Asp
 450 455

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 965 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGCACGAGCT GGACAGCACT ACAAGGCGGG AAGCCATGTC TGT TTTGGAA CGAGACTTTC 60
 35 CAGCATCCAT ACAACACTCT GAAATACCCC AACGGTAGTG GGGGCCTGGG TGAGCACAAAC 120
 TATTGCAGAA ATCCAGATGG AGACGTGAGC CCCTGGTGCT ATGTGGCAGA GCACGAGGAT 180
 GGTGTCTACT GGAAGTACTG TGAGATACCT GCTTGCCAGA TGCTGGAAA CCTTGGCTGC 240
 TACAAGGATC ATGGAAACCC ACCTCCTCTA ACTGGCACCA TAAAAACGTC CAACAACTC 300
 40 ACCATACAAA CTTGCATCAG TTTTGTGCGG AGTCAGAGGT TCAAGTTTGC TGGGATGGAG 360
 TCAGGCTATG CTTGCTTCTG TGGAAACAAT CCGTATTACT GGAAGTACGG GGAGGCAGCC 420
 AGTACCGAAT GCAACAGCGT CTGCTTCGGG GATCACACCC AACCCTGTGG TGGCGATGGC 480
 AGGATCATCC TCTTTGATAC TCTCGTGGGC GCCTGCGGTG GGAAGTACTC AGCCATGTCT 540
 TCTGTGGTCT ATTCCCCTGA CTCCCCGAC ACCTATGCCA CGGGGAGGGT CTGCTACTGG 600
 45 ACCATCCGGG TTCCGGGGGC CTCCCACATC CACTTCAGCT TCCCCCTATT TGACATCAGG 660
 GACTCGGCGG ACATGGTGGA GCTTCTGGAT GGCTACACCC ACCGTGTCCT AGCCCGCTTC 720
 CACGGGAGGA GCCGCCACCC TCTGTCTTTC AACGTCTCTC TGGACTTCGT CATCTTGTAT 780
 TTCTTCTCTG ATCGCATCAA TCAGGCCCAG GGATTGCTG TTTTATACCA AGCCGTCAAG 840
 50 GAAGAAGTGC CACAGGAGAG GCCCGCTGTC AACCAGACGG TGGCCGAGGT GGATCACGGA 900
 GCAGGCCAAC CTTCAGTGTT CAGCGCTTCC CGGTCTCCTC AAAGTCCTAT TATGTGCATG 960
 CACCA 965

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 321 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gly Thr Ser Trp Thr Ala Leu Gln Gly Gly Lys Pro Cys Leu Phe Trp
 1 5 10 15
 Asn Glu Thr Phe Gln His Pro Tyr Asn Thr Leu Lys Tyr Pro Asn Gly
 20 25 30
 Ser Gly Gly Leu Gly Glu His Asn Tyr Cys Arg Asn Pro Asp Gly Asp
 35 40 45
 Val Ser Pro Trp Cys Tyr Val Ala Glu His Glu Asp Gly Val Tyr Trp
 50 55 60
 Lys Tyr Cys Glu Ile Pro Ala Cys Gln Met Pro Gly Asn Leu Gly Cys
 65 70 75 80
 Tyr Lys Asp His Gly Asn Pro Pro Pro Leu Thr Gly Thr Ile Lys Thr
 85 90 95
 Ser Asn Lys Leu Thr Ile Gln Thr Cys Ile Ser Phe Cys Arg Ser Gln
 100 105 110
 Arg Phe Lys Phe Ala Gly Met Glu Ser Gly Tyr Ala Cys Phe Cys Gly
 115 120 125
 Asn Asn Pro Asp Tyr Trp Lys Tyr Gly Glu Ala Ala Ser Thr Glu Cys
 130 135 140
 Asn Ser Val Cys Phe Gly Asp His Thr Gln Pro Cys Gly Gly Asp Gly
 145 150 155 160
 Arg Ile Ile Leu Phe Asp Thr Leu Val Gly Ala Cys Gly Gly Asn Tyr
 165 170 175
 Ser Ala Met Ser Ser Val Val Tyr Ser Pro Asp Phe Pro Asp Thr Tyr
 180 185 190
 Ala Thr Gly Arg Val Cys Tyr Trp Thr Ile Arg Val Pro Gly Ala Ser
 195 200 205
 His Ile His Phe Ser Phe Pro Leu Phe Asp Ile Arg Asp Ser Ala Asp
 210 215 220
 Met Val Glu Leu Leu Asp Gly Tyr Thr His Arg Val Leu Ala Arg Phe
 225 230 235 240

His Gly Arg Ser Arg Pro Pro Leu Ser Phe Asn Val Ser Leu Asp Phe
 245 250 255
 5 Val Ile Leu Tyr Phe Phe Ser Asp Arg Ile Asn Gln Ala Gln Gly Phe
 260 265 270
 Ala Val Leu Tyr Gln Ala Val Lys Glu Glu Leu Pro Gln Glu Arg Pro
 275 280 285
 10 Ala Val Asn Gln Thr Val Ala Glu Val Asp His Gly Ala Gly Gln Pro
 290 295 300
 Ser Val Phe Ser Ala Ser Arg Ser Pro Gln Ser Pro Ile Met Cys Met
 305 310 315 320
 15 His

20 Claims

1. An isolated polypeptide selected from the group consisting of:

25 (i) an isolated polypeptide comprising an amino acid sequence selected from the group having at least:

- (a) 70% identity;
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity

30

to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2;

- (ii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2 or
 (iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2.

35 2. An isolated polynucleotide selected from the group consisting of:

(i) an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least

- (a) 70% identity;
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity;

40

to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ ID NO:2;

45

(ii) an isolated polynucleotide comprising a nucleotide sequence that has at least:

- (a) 70% identity
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity;

50

over its entire length to a nucleotide sequence encoding the polypeptide of SEQ ID NO:2;

(iii) an isolated polynucleotide comprising a nucleotide sequence which has at least:

55

- (a) 70% identity;
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity;

- to that of SEQ ID NO: 1 over the entire length of SEQ ID NO:1;
- (iv) an isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2;
- (v) an isolated polynucleotide which is the polynucleotide of SEQ ID NO: 1; or
- (vi) an isolated polynucleotide obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof;
- or a nucleotide sequence complementary to said isolated polynucleotide.
3. An antibody immunospecific for the polypeptide of claim 1.
4. A method for the treatment of a subject:
- (i) in need of enhanced activity or expression of the polypeptide of claim 1 comprising:
- (a) administering to the subject a therapeutically effective amount of an agonist to said polypeptide; and/or
- (b) providing to the subject an isolated polynucleotide comprising a nucleotide sequence encoding said polypeptide in a form so as to effect production of said polypeptide activity *in vivo*; or
- (ii) having need to inhibit activity or expression of the polypeptide of claim 1 comprising:
- (a) administering to the subject a therapeutically effective amount of an antagonist to said polypeptide; and/or
- (b) administering to the subject a nucleic acid molecule that inhibits the expression of a nucleotide sequence encoding said polypeptide; and/or
- (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said polypeptide for its ligand, substrate, or receptor.
5. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the polypeptide of claim 1 in a subject comprising:
- (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of said polypeptide expression in a sample derived from said subject.
6. A method for screening to identify compounds which stimulate or which inhibit the function of the polypeptide of claim 1 which comprises a method selected from the group consisting of:
- (a) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;
- (b) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;
- (c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes bearing the polypeptide;
- (d) mixing a candidate compound with a solution containing a polypeptide of claim 1, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a standard; or
- (e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide and said polypeptide in cells, using for instance, an ELISA assay.
7. An agonist or an antagonist of the polypeptide of claim 1.
8. An expression system comprising a polynucleotide capable of producing a polypeptide of claim 1 when said expression system is present in a compatible host cell.
9. A host cell comprising the expression system of claim 8 or a membrane thereof expressing the polypeptide of claim 1.
10. A process for producing a polypeptide of claim 1 comprising culturing a host cell of claim 9 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.

11. A process for producing a host cell as defined in claim 9 comprising transforming or transfecting a cell with the expression system of claim 8 such that the host cell, under appropriate culture conditions, produces a polypeptide of claim 1.
12. A recombinant host cell produced by the process of claim 11 or a membrane thereof expressing a polypeptide of claim 1.
13. An isolated polynucleotide selected from the group consisting of:
 - (a) an isolated polynucleotide comprising a nucleotide sequence which has at least 70%, 80%, 90%, 95%, 97% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;
 - (b) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:3;
 - (c) the polynucleotide of SEQ ID NO:3; or
 - (d) an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide which has at least 70%, 80%, 90%, 95%, 97-99% identity to the amino acid sequence of SEQ ID NO:4, over the entire length of SEQ ID NO:4.
14. A polypeptide selected from the group consisting of:
 - (a) a polypeptide which comprises an amino acid sequence which has at least 70%, 80%, 90%, 95%, 97-99% identity to that of SEQ ID NO:4 over the entire length of SEQ ID NO:4;
 - (b) a polypeptide which has an amino acid sequence which is at least 70%, 80%, 90%, 95%, 97-99% identity to the amino acid sequence of SEQ ID NO:4 over the entire length of SEQ ID NO:4;
 - (c) a polypeptide which comprises the amino acid of SEQ ID NO:4;
 - (d) a polypeptide which is the polypeptide of SEQ ID NO:4;
 - (e) a polypeptide which is encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:3.